ES-Certificates save the World - at least Orchard Meadows Commodification of Ecosystem Services



Introduction - What are orchard meadows?

- widespread traditional cultural landscape in Europe (Europe: ca. 1 Mio. ha)
- cultivation of orchard meadows (OM) is officially declared Intangible Cultural Heritage by UNESCO (GER, AU)
- traditional agroforestry system: OM combines fruit tree growing most frequently with permanent pasture.
 - large-crowned fruit trees in wide distances, mostly high-trunked and strong-growing
 - o characterized by different age, size, as well as fruit varieties, and species (Herzog and Oetmann 2001; Herzog 1998)

Current status - Traditional cultural landscape with very high biodiversity and extent of ES

- traditional agroforestry system with the "greatest" biodiversity and extent of ecosystem services of all cultural landscapes: 5,000 animal and plant species (Rösler 1993)
- decline in stocks of 60-70% since 1950: threatened with complete destruction (red list of endangered biotope types in Germany) (BMUV, 2017)
- lack of economic basis due to the considerable amount of time and effort required for maintenance and preservation compared to income from product sales
- □ In line with other traditional systems, long term conservation of the OM system needs financial incentives for farmers, either by market or by public subsidies (Flinzberger et al. 2020; Forejt and Syrbe 2019; Herzog 1998; Plieninger et al. 2013; Schaich et al. 2010; Tojnko et al. 2011; Wolpert et al. 2020).


Examples for special ecosystem services of OM

- 👲 food: fresh & dried fruit, fruit juice & juice spritzer, jam, cider/perry, Secco
- onimal food: gras/hey/silage, fruit, tree foliage (extension of the grazing period)
- carbon sequestration through permanent grassland and trees
- climate regulation: microclimate cold air production
- gene reservoir: 5.000+ fruit types (climate-resistant varieties)
- Intangible Cultural Heritage & identification: influence on society (customs, festivals, traditions), regional identity
- health effect: landscape lowers blood pressure, old fruit varieties suitable for allergy sufferers, hyposensitization for allergy sufferers



Type of landscape	potential total species occurrence
arable land (wheat and maize)	6,54
modern agroforestry system (walnut trees for timber produc- tion)	7,98
orchard meadows (high-stemmed cherry trees)	13,10

Monitoring of species occurrence in different agroforestry systems. Orchard meadows with the highest species occurrence (source: own illustration based von Kay et al. (2019))





ES certificates - Rewarding the provision and conservation of biodiversity and ecosystem services

Aim: Integration of ecosystems services in value chains via ES-Certificates and preserving high levels of biodiversity and ES

Current status


- lack of profitability ⇒ termination of management (cultivation)
- decreasing extent of biodiversity and ES due to missing replacement planting, loss of trees, lack of underutilisation, succession, etc.

Short term scenario (with ES-Certificates)


- commodification of ES via certificates (and other paths: more funding, higher sales revenues)
 - ⇒ economic efficiency given
 - ⇒ preserving of biodiversity and ES

Long term scenario (ES-certificates included cultivation guideline)


- mandatory cultivation guideline for certified OM areas
- consulting & support for further development of certified OM areas
 - ⇒ increase in biodiversity and scope of ESS



Schematic value chain orchard meadows with integration of biodiversity and ES (source: own illustration)



Schematic description of the implementation of ES certificate in practice (source: own illustration).



References

Bundesministerium für Umwelt, Naturschutz, Bau und Reaktorsicherheit (BMUV) (2017): Rote Liste gefährdeter Biotoptypen. Bundespressekonferenz am 31. Mai 2017.

Forejt M, Syrbe RU (2019) The current status of orchard meadows in Central Europe: multi-source area estimation in Saxony (Germany) and the Czech Republic. Moravian Geographical Reports 27(4): 217–228. https://doi.org/10.2478/mgr-2019-0017

Flinzberger L, Zinngrebe Y, Plieninger T (2020). Labelling in Mediterranean agroforestry landscapes: a Delphi study on relevant sustainability indicators. Sustainability Science 15:1369–1382. https://doi.org/https://doi.org/10.1007/s11625-020-00800-2

Herzog F., Oetmann A (2001) Communities of interest and agroecosystem restoration: Streuobst in Europe. In: Flora C (ed) Interactions Between Agroecosystems and Rural Communities, CRC Press, pp. 85–99. https://doi.org/10.1201/9781420041385.ch7

Herzog F (1998) Streuobst: a traditional agroforestry system as a model for agroforestry development in temperate Europe. Agroforestry Systems 42:61–80.

Kay, S.; Jeanneret, P.; Herzog, F. (2019): Umweltleistungen: Traditionelle Agroforstsysteme sind Biodiversitäts-Hotspot – gilt dies auch für moderne Systeme? Agroscope. Forschungsgruppe Agrarlandschaft und Biodiversität, Schweiz.

Rösler, M. (1993): Streuobstbau - Behauptungen und Realitäten - Einführung in die Tagung; Materialien 5/93 der Ev. Akademie Bad Boll -Vielfalt in aller Munde - Perspektiven für Bewirtschaftung und Vermarktung im bundesweiten Streuobstbau; Dokumentation der Tagung von Ev. Akademie Bad Boll und NABU-Bundesverband; Hrsg. Markus Rösler & Jobst Krauß; S. 5-10

Schaich H, Bieling C. Plieninger T (2010) Linking ecosystem services with cultural landscape research. GAIA-Ecological Perspectives for Science and Society 19(4):269– 277. https://doi.org/10.14512/gaia.19.4.9

Tojnko S, Rozman Č, Unuk T, Pažek K, Pamič S (2011) A qualitative multi-attribute model for the multifunctional assessment of "Streuobst stands" in NE Slovenia. Erwerbs-Obstbau 53(4):157–166. https://doi.org/10.1007/s10341-011-0149-0

Wolpert F, Quintas-Soriano C, Plieninger T (2020) Exploring land-use histories of treecrop landscapes: a cross-site comparison in the Mediterranean Basin. Sustainability Science 15(5):1267–1283. https://doi.org/10.1007/s11625-020-00806-w

